

l17_arytm_1

(TMUxbjjn6kCjNxuZvZWp9y1JoCwhKyC8icH)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_arytm_2 : \iota$ be given. Let $k11_arytm_3 : \iota$ be given. Let $k1_arytm_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_arytm_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_arytm_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_arytm_2 : \iota \Rightarrow \iota$ be given. Let $k5_arytm_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_arytm_2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 X1 k2_arytm_2) \Rightarrow ((X0 = k11_arytm_3) \Rightarrow (r1_arytm_2 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 X1 k2_arytm_2) \Rightarrow (((X1 = k11_arytm_3) \Rightarrow (k7_arytm_2 X0 X1 = X0)) \wedge ((X0 = k11_arytm_3) \Rightarrow (k7_arytm_2 X0 X1 = X1)) \wedge (\neg(X1 \neq k11_arytm_3) \wedge ((X0 \neq k11_arytm_3) \wedge (k7_arytm_2 X0 X1 \neq k4_arytm_2 (k5_arytm_2 (k3_arytm_2 X0) (k3_arytm_2 X1)))))))) \quad (2)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 X1 k2_arytm_2) \Rightarrow (\forall X2.(m1_subset_1 X2 k2_arytm_2) \Rightarrow (((r1_arytm_2 X1 X0) \Rightarrow ((X2 = k1_arytm_1 X0 X1) \Leftrightarrow (k7_arytm_2 X2 X1 = X0))) \wedge ((\neg r1_arytm_2 X1 X0) \Rightarrow ((X2 = k1_arytm_1 X0 X1) \Leftrightarrow (X2 = k11_arytm_3)))))) \quad (3)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 X1 k2_arytm_2) \Rightarrow ((X0 = k11_arytm_3) \Rightarrow (k1_arytm_1 X1 X0 = X1)))$$