

l17_polyeq_1 (TMQco- hAyg4CPCrjfv5LoTRrQaZWR AoHyJV)

October 27, 2020

Let $v1_abian : \iota \Rightarrow o$ be given. Let $np_3 : \iota$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $r1_int_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & ((v2_xreal_0 \ np_2) \wedge (m2_subset_1 \ np_2 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_2 \ k5_numbers) \wedge (m1_subset_1 \ np_2 \ k1_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$k2_xcmplx_0 \ np_2 \ np_1 = np_3 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_int_1 \ X0) \wedge (v1_int_1 \ X1)) \Rightarrow (r1_int_1 \ X0 \ X0) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. ((v1_int_1 \ X0) \wedge (v1_abian \ X0)) \Rightarrow (\neg v1_abian \ (k2_xcmplx_0 \ X0 \ np_1)) \quad (5)$$

Assume the following.

$$\forall X0. (v1_int_1 \ X0) \Rightarrow ((v1_abian \ X0) \Leftrightarrow (r1_int_1 \ np_2 \ X0)) \quad (6)$$

Assume the following.

$$\forall X0. (m1_subset_1 \ X0 \ k4_ordinal1) \Rightarrow (v7_ordinal1 \ X0) \quad (7)$$

Assume the following.

$$\forall X0. (v7_ordinal1 \ X0) \Rightarrow (v1_int_1 \ X0) \quad (8)$$

Theorem 1 $\neg v1_abian \ np_3$.