

1183_group_9 (TMU-
vwMtF6ryTNc8LXfmZAFnv1Y55yBtymRW)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k2_xcmplx_0 X0 X1) X2) \Leftrightarrow (r1_xxreal_0 \\ & X0 (k6_xcmplx_0 X2 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 (k6_xcmplx_0 X0 np_1)))) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_subset_1\ X1\ k5_numbers))\Rightarrow (k1_nat_1\ X0\ X1 = k2_xcmplx_0\ X0\ X1) \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0\ k4_ordinal1)\wedge(v3_ordinal1\ k4_ordinal1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0\ X0)\wedge((\neg v1_xboole_0\ X1)\wedge (m1_subset_1\ X1\ (k1_zfmisc_1\ X0))))\Rightarrow(\forall X2.(m2_subset_1\ X2\ X0\ X1)\Rightarrow(m1_subset_1\ X2\ X0)) \quad (9)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (11)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xreal_0\ X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(v1_xreal_0\ X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0))\Rightarrow(v1_xboole_0\ X1)) \quad (14)$$

Theorem 1

$$\forall X0.(m2_subset_1\ X0\ k1_numbers\ k5_numbers)\Rightarrow(\forall X1.(m2_subset_1\ X1\ k1_numbers\ k5_numbers)\Rightarrow((\neg r1_xxreal_0\ X1\ X0)\Leftrightarrow (r1_xxreal_0\ X0\ (k6_xcmplx_0\ X1\ np_1))))$$