

118_xreal_1 (TM-
PqHc5fvoEHsZ9SxrCw2KTxHXWmvfCsxk3)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_0 : \iota$ be given. Let $k11_arytm_3 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 k6_numbers = X0) \quad (3)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$k11_arytm_3 = k1_xboole_0 \quad (6)$$

Assume the following.

$$\exists X0. (v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (7)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 X0) \Rightarrow (\forall X1. (v1_xcmplx_0 X1) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (X1 = k7_xcmplx_0 (k3_xcmplx_0 X1 X0) X0))) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1))) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 (k4_xcmplx_0 X1) (k4_xcmplx_0 X0)))) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 k6_numbers X2)) \Rightarrow (r1_xxreal_0 (k3_xcmplx_0 X0 X2) (k3_xcmplx_0 X1 X2)))))) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k4_xcmplx_0 (k4_xcmplx_0 X0) = X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k3_xcmplx_0 X0 X1)) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow ((X1 = k4_xcmplx_0 X0) \Leftrightarrow (k2_xcmplx_0 X0 X1 = k6_numbers))) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow ((r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (16)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (18)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 X2 X1) \wedge (r1_xxreal_0 (k3_xcmplx_0 X2 X0) (k3_xcmplx_0 X1 X0)))))))$$