

## l1\_euler\_2

(TMUd2ma8pQJhRJawEuRXAuyvmqrPG8qv7j9)

October 27, 2020

Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v3\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((r1\_xreal\_0 X0 X1) \Rightarrow ((v1\_xboole\_0 X0) \vee ((v2\_xreal\_0 X1) \vee (v3\_xreal\_0 X0)))))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (2)$$

Assume the following.

$$((v2\_xreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (3)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (4)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0) \wedge ((v1\_xcmplx\_0 X0) \wedge ((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X0))) \quad (5)$$

Assume the following.

$$\forall X0.(v1\_int\_1 X0) \Rightarrow ((\neg r1\_xreal\_0 X0 k6\_numbers) \Rightarrow (r1\_xreal\_0 np\_1 X0)) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xreal\_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.((v1\_xxreal\_0 X0)\wedge(v2\_xxreal\_0 X0))\Rightarrow((\neg v1\_xboole\_0 X0)\wedge((v1\_xxreal\_0 X0)\wedge(\neg v3\_xxreal\_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.(v1\_int\_1 X0)\Rightarrow(v1\_xreal\_0 X0) \quad (9)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers)\Rightarrow(\neg v3\_xxreal\_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (11)$$

**Theorem 1**

$$\forall X0.(v1\_int\_1 X0)\Rightarrow((\neg r1\_xxreal\_0 np\_1 X0)\Leftrightarrow(r1\_xxreal\_0 X0 k1\_xboole\_0))$$