

# l1\_integr11

(TMXzyHUSy5ineX7twSvJxdEV33Y3s2WEtJD)

October 27, 2020

Let  $k2\_subset\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_fcont\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k8\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k32\_sin\_cos : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow(k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (6)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0)\wedge((v1\_xcmplx\_0 X0)\wedge((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X0))) \quad (7)$$

Assume the following.

$$k1\_relset\_1 k1\_numbers (k1\_fcont\_1 (k10\_real\_1 np\_1 (k8\_real\_1 np\_2 k32\_sin\_cos)) k6\_numbers) = k1\_numbers \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers)\wedge(v1\_xreal\_0 X1))\Rightarrow(m1\_subset\_1 (k8\_real\_1 X0 X1) k1\_numbers) \quad (9)$$

Assume the following.

$$m1\_subset\_1 k32\_sin\_cos k1\_numbers \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow((v1\_funct\_1 (k1\_fcont\_1 X0 X1))\wedge((v1\_funct\_2 (k1\_fcont\_1 X0 X1) k1\_numbers k1\_numbers)\wedge(m1\_subset\_1 (k1\_fcont\_1 X0 X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_numbers k1\_numbers)))))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers)\wedge(v1\_xreal\_0 X1))\Rightarrow(m1\_subset\_1 (k10\_real\_1 X0 X1) k1\_numbers) \quad (12)$$

Assume the following.

$$\forall X0.k2\_subset\_1 X0 = X0 \quad (13)$$

Assume the following.

$$k1\_xboole\_0 = the (\lambda X0 : \iota.v1\_xboole\_0 X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(((X1\neq k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 X1)\Leftrightarrow(X0 = k1\_relset\_1 X0 X2)))\wedge((X1 = k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 X1)\Leftrightarrow(X2 = k1\_xboole\_0)))) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (16)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (18)$$

**Theorem 1**

$$k2\_subset\_1 k1\_numbers = k9\_xtuple\_0 (k1\_fcont\_1 (k10\_real\_1 np\_1 np\_2) k6\_numbers)$$