

l1_jordan1c (TMKV- TuybW1n75wmLUnrmvvqEh6rpoo5XwLv)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_int_1 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_int_1 X1) \Rightarrow (k2_xcmplx_0 (k1_int_1 X0) X1 = k1_int_1 (k2_xcmplx_0 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (X0 = k6_xcmplx_0 (k2_xcmplx_0 X0 X1) X1)) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_int_1 (k1_int_1 X0)) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (v1_xreal_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_int_1 X0) \quad (9)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow (k6_xcmplx_0 (k1_int_1 (k2_xcmplx_0 X0 X1)) X1 = k1_int_1 X0))$$