

## l1\_limfunc2

(TMVJbY5cHU3cKhK2CBarG566akdASwaQezw)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X1 X2) \Rightarrow ((r1\_xxreal\_0 X0 k6\_numbers) \vee \\ & ((\neg r1\_xxreal\_0 X2 (k6\_xcmplx\_0 X1 X0)) \wedge (\neg r1\_xxreal\_0 (k2\_xcmplx\_0 \\ & X2 X0) X1)))))) \end{aligned} \tag{1}$$

### Theorem 1

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 X2) \Rightarrow ((r1\_xxreal\_0 X1 k6\_numbers) \vee \\ & ((\neg r1\_xxreal\_0 X2 (k6\_xcmplx\_0 X0 X1)) \wedge (\neg r1\_xxreal\_0 (k2\_xcmplx\_0 \\ & X2 X1) X0)))))) \end{aligned}$$