

l1_pdiff_7

(TMK9QhPbwAcLkYSaowUVXgeGAkbixpi8gJZ)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_normsp_0 : \iota \Rightarrow o$ be given. Let $v4_normsp_0 : \iota \Rightarrow o$ be given. Let $v2_normsp_1 : \iota \Rightarrow o$ be given. Let $l1_normsp_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_nfcont_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_normsp_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\ & (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_nfcont_1 X2 X0 X1) \Rightarrow (\forall X3. (m1_nfcont_1 X3 X0 X1) \Rightarrow (\exists X4. (m1_nfcont_1 X4 X0 X1) \wedge ((r1_tarski X4 X2) \wedge (r1_tarski X4 X3))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X0 X2)) \Rightarrow (r1_tarski X0 (k3_xboole_0 X1 X2)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge \\ & ((v2_rlvect_1 X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 \\ & X0) \wedge ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge \\ & ((v3_normsp_0 X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 \\ & X0)))))))))) \wedge (m1_subset_1 X1 (u1_struct_0 X0))) \Rightarrow (\forall X2. \\ & (m1_nfcont_1 X2 X0 X1) \Rightarrow (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 \\ & X0)))) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (m1_subset_1 (k9_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0. (((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 \\ & X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\ & (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_subset_1 \\ & X2 (k1_zfmisc_1 (u1_struct_0 X0)) \Rightarrow ((m1_nfcont_1 X2 X0 X1) \Leftrightarrow (\exists X3. \\ & (m1_subset_1 X3 k1_numbers) \wedge ((\neg r1_xxreal_0 X3 k6_numbers) \wedge \\ & r1_tarski (ReplSep (toset (\lambda X4 : \iota. m1_subset_1 X4 (u1_struct_0 \\ & X0))) (\lambda X4 : \iota. \neg r1_xxreal_0 X3 (k1_normsp_0 X0 (k5_algstr_0 \\ & X0 X4 X1))) (\lambda X4 : \iota. X4) X2)))))) \end{aligned} \tag{7}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k9_subset_1 X0 X2 X1) \tag{8}$$

Theorem 1

$$\begin{aligned} & \forall X0. (((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 \\ & X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\ & (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_nfcont_1 \\ & X2 X0 X1) \Rightarrow (\forall X3. (m1_nfcont_1 X3 X0 X1) \Rightarrow (m1_nfcont_1 (k9_subset_1 \\ & (u1_struct_0 X0) X2 X3) X0 X1)))) \end{aligned}$$