

ll_trees_9

(TMT3XogXvU2zcyjURTYN4DM7ZjLU1YU5yMe)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow (\neg (r1_xxreal_0 X0 X1) \wedge (\forall X2. (v7_ordinal1 X2) \Rightarrow (X1 \neq k2_xcmplx_0 X0 X2)))) \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 \ np_1) \wedge (m2_subset_1 \ np_1 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_1 \ k5_numbers) \wedge (m1_subset_1 \ np_1 \ k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 \ X0) \wedge ((\neg v1_xboole_0 \ X1) \wedge \\ & (m1_subset_1 \ X1 \ (k1_zfmisc_1 \ X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & \ X2 \ X0 \ X1) \Leftrightarrow (m1_subset_1 \ X2 \ X1)) \end{aligned} \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow \\ & (k4_finseq_1 \ X0 = k9_xtuple_0 \ X0) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((m1_subset_1 \ X0 \ k5_numbers) \wedge (v7_ordinal1 \\ & \ X1)) \Rightarrow (k2_nat_1 \ X0 \ X1 = k2_xcmplx_0 \ X0 \ X1) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 \ X1) \wedge (v4_relat_1 \ X1 \ X0)) \Rightarrow (\\ & \ k1_relset_1 \ X0 \ X1 = k9_xtuple_0 \ X1) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v7_ordinal1 \ X0) \wedge (m1_subset_1 \ X1 \ k5_numbers)) \Rightarrow \\ & (k1_nat_1 \ X0 \ X1 = k2_xcmplx_0 \ X0 \ X1) \end{aligned} \quad (12)$$

Assume the following.

$$(\neg v1_xboole_0 \ k4_ordinal1) \wedge (v3_ordinal1 \ k4_ordinal1) \quad (13)$$

Assume the following.

$$m1_subset_1 \ k5_numbers \ (k1_zfmisc_1 \ k1_numbers) \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow \\ & (m1_subset_1 \ (k4_finseq_1 \ X0) \ (k1_zfmisc_1 \ k5_numbers)) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow \\ & (m2_subset_1 \ (k3_finseq_1 \ X0) \ k1_numbers \ k5_numbers) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Leftrightarrow(X0 \in k4_ordinal1) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1\ X0\ k5_numbers)\wedge(v7_ordinal1\ X1))\Rightarrow(k2_nat_1\ X0\ X1 = k2_nat_1\ X1\ X0) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_subset_1\ X1\ k5_numbers))\Rightarrow(k1_nat_1\ X0\ X1 = k1_nat_1\ X1\ X0) \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (20)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow((v1_relat_1\ X0)\wedge((v4_relat_1\ X0\ k5_numbers)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))) \quad (21)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0))\Rightarrow(v1_xboole_0\ X1)) \quad (22)$$

Theorem 1

$$\forall X0.\forall X1.((v1_relat_1\ X1)\wedge((v1_funct_1\ X1)\wedge(v1_finseq_1\ X1)))\Rightarrow(\neg(X0 \in k4_finseq_1\ X1)\wedge(\forall X2.(m1_subset_1\ X2\ k5_numbers)\Rightarrow(\neg(X0 = k2_nat_1\ X2\ np_1)\wedge(\neg r1_xreal_0\ (k3_finseq_1\ X1\ X2))))))$$