

l204_xreal_1

(TMT885zZwQsjPw9iCGJDRZHK8aquMimXNh)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k11_arytm_3 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 \\ (k5_xcmplx_0 X0) (k5_xcmplx_0 X1)) \Rightarrow ((r1_xxreal_0 (k5_xcmplx_0 \\ X0) k6_numbers) \vee (r1_xxreal_0 X1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (3)$$

Assume the following.

$$k11_arytm_3 = k1_xboole_0 \quad (4)$$

Assume the following.

$$k5_xcmplx_0 np_1 = np_1 \quad (5)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow ((v2_xxreal_0 X0) \Leftrightarrow (\neg r1_xxreal_0 \\ X0 k6_numbers)) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg(\neg r1_xxreal_0 X0 \text{ np--1}) \wedge (r1_xxreal_0 \text{ np--1} (k5_xcmplx_0 X0)))$$