

l207\_xreal\_1  
(TMF115ogeLvS1fsPQSzyznEg1rhdxWYoXiM)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow ((X0 \neq k6\_numbers) \Rightarrow (k7\_xcmplx\_0 X0 X0 = np\_1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k7\_xcmplx\_0 k6\_numbers X0 = k6\_numbers) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 X0 X2)))) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 (k7\_xcmplx\_0 X1 X0) (k7\_xcmplx\_0 X2 X0)))))) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 k6\_numbers X0)) \Rightarrow (r1\_xxreal\_0 (k7\_xcmplx\_0 X0 X1) np\_1))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (v1\_xreal\_0 (k7\_xcmplx\_0 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (7)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 k6\_numbers X1)) \Rightarrow (r1\_xxreal\_0 (k7\_xcmplx\_0 X0 X1) np\_1)))$$