

l20_complex1

(TMGmhJsp42pDMZnaDrKsWRT8zpzFeMusUJp)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k3_arytm_0 : \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_arytm_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 (k4_xcmplx_0 X0) (k4_xcmplx_0 X1) = k4_xcmplx_0 (k2_xcmplx_0 X0 X1)) \quad (2)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k1_real_1 X0 = k4_xcmplx_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((X0 = X2) \wedge (X1 = X3)) \Rightarrow (k1_arytm_0 X0 X1 = k2_xcmplx_0 X2 X3)))))) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k4_xcmplx_0 (k4_xcmplx_0 X0) = X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(v1_xreal_0 (k4_xcmplx_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.((\neg v3_xxreal_0 X0)\wedge(v1_xreal_0 X0))\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(\neg v2_xxreal_0 (k4_xcmplx_0 X0))) \quad (9)$$

Assume the following.

$$\forall X0.((\neg v2_xxreal_0 X0)\wedge(v1_xreal_0 X0))\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(\neg v3_xxreal_0 (k4_xcmplx_0 X0))) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(\forall X1.(m1_subset_1 X1 k1_numbers)\Rightarrow((X1 = k3_arytm_0 X0)\Leftrightarrow(k1_arytm_0 X0 X1 = k6_numbers))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0)\wedge(v1_xcmplx_0 X1))\Rightarrow(k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (13)$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0)\wedge((v1_xxreal_0 X0)\wedge(\neg v2_xxreal_0 X0)))\Rightarrow((v1_xxreal_0 X0)\wedge(v3_xxreal_0 X0)) \quad (14)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (15)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0)\wedge(v2_xxreal_0 X0))\Rightarrow((\neg v1_xboole_0 X0)\wedge((v1_xxreal_0 X0)\wedge(\neg v3_xxreal_0 X0))) \quad (16)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (17)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xcmplx_0 X0) \quad (18)$$

Theorem 1 $\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(k3_arytm_0 X0 = k1_real_1 X0)$.