

## l21\_anproj\_2

(TMTYxK1Z9vTMHrqPf48TqjGy49azKUjPQc1)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k1\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v9\_struct\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $l2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_algstr\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge ( \\ (v13\_algstr\_0 X1) \wedge ((v2\_rlvect\_1 X1) \wedge ((v3\_rlvect\_1 X1) \wedge ((v4\_rlvect\_1 \\ X1) \wedge ((v5\_rlvect\_1 X1) \wedge ((v6\_rlvect\_1 X1) \wedge ((v7\_rlvect\_1 X1) \wedge \\ ((v8\_rlvect\_1 X1) \wedge (l1\_rlvect\_1 X1)))))))))) \Rightarrow (\forall X2.(m1\_subset\_1 \\ X2 (u1\_struct\_0 X1)) \Rightarrow (\neg(k1\_rlvect\_1 X1 X2 X0 = k4\_struct\_0 X1) \wedge \\ ((X0 \neq k6\_numbers) \wedge (X2 \neq k4\_struct\_0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(l2\_algstr\_0 X0) \Rightarrow ((l2\_struct\_0 X0) \wedge (l1\_algstr\_0 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(l1\_rlvect\_1 X0) \Rightarrow (l2\_algstr\_0 X0) \quad (3)$$

Assume the following.

$$\forall X0.(l2\_struct\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ X0)) \Rightarrow ((v9\_struct\_0 X1 X0) \Leftrightarrow (X1 = k4\_struct\_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (5)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge (v13\_algstr\_0 X0) \wedge (v2\_rlvect\_1 \\ & X0) \wedge (v3\_rlvect\_1 X0) \wedge (v4\_rlvect\_1 X0) \wedge (v5\_rlvect\_1 X0) \wedge \\ & ((v6\_rlvect\_1 X0) \wedge (v7\_rlvect\_1 X0) \wedge (v8\_rlvect\_1 X0) \wedge (l1\_rlvect\_1 \\ & X0)))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow \\ & (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3.(m1\_subset\_1 \\ X3 k1\_numbers) \Rightarrow (\neg(k1\_rlvect\_1 X0 X1 X3 = X2) \wedge (X3 \neq k6\_numbers) \wedge \\ & ((\neg v9\_struct\_0 X1 X0) \wedge (v9\_struct\_0 X2 X0)))))) \end{aligned}$$