

## l22\_compos\_2

(TMV8kKrW4ZRR5L1KAFkm8st4j28PHiQzbfaf)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k6\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k16\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. k16\_funcop\_1 X0 X1 = k1\_tarski (k4\_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 X1 X0 \in k10\_xtuple\_0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (r1\_tarski X0 X1) \Rightarrow ((X2 \in X0) \vee (r1\_tarski X0 (k4\_xboole\_0 X1 (k1\_tarski X2)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((k4\_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9\_xtuple\_0 X2) \wedge (X1 = k1\_funct\_1 X2 X0))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k6\_subset\_1 X0 X1 = k4\_xboole\_0 X0 X1 \quad (5)$$

### Theorem 1

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (\forall X2. \forall X3. (\neg X2 \in k10\_xtuple\_0 X0) \Rightarrow (r1\_tarski X0 (k6\_subset\_1 X1 (k16\_funcop\_1 X3 X2)))))))$$