

l22_jordan1b (TM-
FRZ1vYB4umHeXL4Phx3ocgQieb1QeRV2M)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k7_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k7_nat_d (k2_xcmplx_0 X0 X1) X1 = X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_subset_1\ X1\ k5_numbers))\Rightarrow (k1_nat_1\ X0\ X1 = k2_xcmplx_0\ X0\ X1) \quad (7)$$

Assume the following.

$$v6_membered\ k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(v7_ordinal1\ X1))\Rightarrow(v7_ordinal1\ (k2_xcmplx_0\ X0\ X1)) \quad (9)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xxreal_0\ X0) \quad (10)$$

Assume the following.

$$\forall X0.(v6_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v7_ordinal1\ X1)) \quad (11)$$

Theorem 1

$$\forall X0.(m1_subset_1\ X0\ k5_numbers)\Rightarrow(\forall X1.(m1_subset_1\ X1\ k5_numbers)\Rightarrow(\neg(r1_xxreal_0\ X0\ X1)\wedge((r1_xxreal_0\ X1\ (k2_nat_1\ X0\ np_1))\wedge((X0\neq X1)\wedge(X0\neq k7_nat_d\ X1\ np_1))))))$$