

l22_mesfun6c

(TMRchDq9PXt4nUg6rQXMw9AJJXfJ8gNx86s)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_mesfunc2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
 & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
 & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
 & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
 & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
 & (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & X0 k7_numbers)))) \Rightarrow ((v6_supinf_2 (k1_mesfunc2 X0 X3) \wedge ((v6_supinf_2 \\
 & (k2_mesfunc2 X0 X3) \wedge (v6_supinf_2 (k10_mesfunc1 X0 X3)))))) \\
 & \tag{1}
 \end{aligned}$$

Theorem 1

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
 & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
 & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
 & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
 & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
 & (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & X0 k7_numbers)))) \Rightarrow ((v6_supinf_2 (k1_mesfunc2 X0 X3) \wedge (v6_supinf_2 \\
 & (k2_mesfunc2 X0 X3))))))
 \end{aligned}$$