

l22_modelc_2
(TMdB1gx51dUh2usYeVvwCx9hScTzTnfumoK)

October 27, 2020

Let $v1_modelc_2 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v3_modelc_2 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k3_modelc_2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow (k1_funct_1 (k7_finseq_1 (k9_finseq_1 X0) X1) np_1 = X0) \quad (1)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_finseq_1 X1 X0) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (5)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (8)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow (m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (11)$$

Assume the following.

$$\forall X0.(m2_finseq_1 X0 k5_numbers)\Rightarrow(k3_modelc_2 X0 = k8_finseq_1 k5_numbers (k12_finseq_1 k5_numbers k6_numbers) X0) \quad (12)$$

Assume the following.

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v3_modelc_2 X0)\Leftrightarrow(\exists X1.((v1_modelc_2 X1)\wedge(m2_finseq_1 X1 k5_numbers))\wedge(X0 = k3_modelc_2 X1))) \quad (13)$$

Theorem 1

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v3_modelc_2 X0)\Rightarrow(k1_funct_1 X0 np_1 = k6_numbers))$$