

l23_sincos10 (TMWxu-
vkMhYNoQ7Wd8GMaiPHyqDvWvoSz85U)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k32_sin_cos : \iota$ be given. Let $np_2 : \iota$ be given. Let $k4_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k31_sin_cos : \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (r1_tarski (k4_xxreal_1 X0 X1) (k3_xxreal_1 X0 X1))) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (k4_rcomp_1 X0 X1 = k3_xxreal_1 X0 X1) \quad (3)$$

Assume the following.

$$k32_sin_cos = k31_sin_cos \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k2_rcomp_1 X0 X1 = k4_xxreal_1 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k10_real_1 X0 X1 = k7_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k7_xcmplx_0 X0 X1)) \quad (7)$$

Assume the following.

$$m1_subset_1 k32_sin_cos k1_numbers \quad (8)$$

Assume the following.

$$v1_xreal_0 k31_sin_cos \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$r1_tarski (k2_rcomp_1 (k10_real_1 k32_sin_cos np_2) k32_sin_cos) \\ (k4_rcomp_1 (k10_real_1 k32_sin_cos np_2) k32_sin_cos)$$