

l24_sincos10 (TMSdnpsaEXJXMT- dTn6bZEntHhE2LaEtnJE9)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k32_sin_cos : \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_fdiff_9 : \iota$ be given. Let $k4_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$r1_tarski (k4_rcomp_1 (k10_real_1 k32_sin_cos np_2) k32_sin_cos) \\ (k1_relset_1 k1_numbers k1_fdiff_9) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski \\ X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (2)$$

Assume the following.

$$r1_tarski (k2_rcomp_1 (k10_real_1 k32_sin_cos np_2) k32_sin_cos) \\ (k4_rcomp_1 (k10_real_1 k32_sin_cos np_2) k32_sin_cos) \quad (3)$$

Theorem 1

$$r1_tarski (k2_rcomp_1 (k10_real_1 k32_sin_cos np_2) k32_sin_cos) \\ (k1_relset_1 k1_numbers k1_fdiff_9)$$