

l25_fib_num4
(TMUTxtd55AV2yVffEELbDpion5zCwVuvJBi)

October 27, 2020

Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $np_5 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_square_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge (r1_xxreal_0 (k6_square_1 X0) k6_numbers)) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_5) \wedge (m2_subset_1 np_5 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_5 k5_numbers) \wedge (m1_subset_1 np_5 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (4)$$

Assume the following.

$$\neg r1_xxreal_0 np_5 np_0 \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k7_square_1 X0 = k6_square_1 X0) \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1 $\neg r1_xxreal_0 (k7_square_1 np_5) k6_numbers.$