

# l25\_goboard7 (TMY- HHA1qmJbBdrdpUGM57oudpZQsrAaWak5)

October 27, 2020

Let  $k9\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k10\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v3\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0\ np\_2) \wedge (m2\_subset\_1\ np\_2\ k1\_numbers\ k5\_numbers)) \wedge \\ & ((m1\_subset\_1\ np\_2\ k5\_numbers) \wedge (m1\_subset\_1\ np\_2\ k1\_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0\ np\_1) \wedge (m2\_subset\_1\ np\_1\ k1\_numbers\ k5\_numbers)) \wedge \\ & ((m1\_subset\_1\ np\_1\ k5\_numbers) \wedge (m1\_subset\_1\ np\_1\ k1\_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$k6\_xcmplx\_0\ np\_1\ (k7\_xcmplx\_0\ np\_1\ np\_2) = k7\_xcmplx\_0\ np\_1\ np\_2 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1\ X0\ k1\_numbers) \wedge (v1\_xreal\_0\ X1)) \Rightarrow (k9\_real\_1\ X0\ X1 = k6\_xcmplx\_0\ X0\ X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1\ X0\ k1\_numbers) \wedge (v1\_xreal\_0\ X1)) \Rightarrow (k10\_real\_1\ X0\ X1 = k7\_xcmplx\_0\ X0\ X1) \quad (5)$$

Assume the following.

$$v3\_membered\ k1\_numbers \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1\ X0\ k1\_numbers) \wedge (v1\_xreal\_0\ X1)) \Rightarrow (m1\_subset\_1\ (k10\_real\_1\ X0\ X1)\ k1\_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(v3\_membered\ X0) \Rightarrow (\forall X1.(m1\_subset\_1\ X1\ X0) \Rightarrow (v1\_xreal\_0\ X1)) \quad (8)$$

**Theorem 1**  $k9\_real\_1\ np\_1\ (k10\_real\_1\ np\_1\ np\_2) = k10\_real\_1\ np\_1\ np\_2$ .