

## l26\_modelc\_1

(TMHV4GXwBj6eM1LfHySaSPZcHsXFnvJ7H9f)

October 27, 2020

Let  $v1\_modelc\_1 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v4\_modelc\_1 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_modelc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (k1\_funct\_1 (k7\_finseq\_1 (k9\_finseq\_1 X0) X1) np\_1 = X0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow \\ (\forall X2. ((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2)))) \Rightarrow \\ (k7\_finseq\_1 (k7\_finseq\_1 X0 X1) X2 = k7\_finseq\_1 X0 (k7\_finseq\_1 X1 X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0. k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1\_finseq\_1 X1 X0)\wedge(m1\_finseq\_1 X2 X0))\Rightarrow(k8\_finseq\_1 X0 X1 X2 = k7\_finseq\_1 X1 X2) \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow(k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (8)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \quad (11)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k9\_finseq\_1 X0))\wedge(v1\_funct\_1 (k9\_finseq\_1 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1\_finseq\_1 X1 X0)\wedge(m1\_finseq\_1 X2 X0))\Rightarrow(m2\_finseq\_1 (k8\_finseq\_1 X0 X1 X2) X0) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow(m2\_finseq\_1 (k12\_finseq\_1 X0 X1) X0) \quad (14)$$

Assume the following.

$$\forall X0.(m2\_finseq\_1 X0 k5\_numbers)\Rightarrow(\forall X1.(m2\_finseq\_1 X1 k5\_numbers)\Rightarrow(k8\_modelc\_1 X0 X1 = k8\_finseq\_1 k5\_numbers (k8\_finseq\_1 k5\_numbers (k12\_finseq\_1 k5\_numbers np\_1) X0) X1)) \quad (15)$$

Assume the following.

$$\forall X0.((v1\_modelc\_1 X0)\wedge(m2\_finseq\_1 X0 k5\_numbers))\Rightarrow((v4\_modelc\_1 X0)\Leftrightarrow(\exists X1.((v1\_modelc\_1 X1)\wedge(m2\_finseq\_1 X1 k5\_numbers))\wedge(\exists X2.((v1\_modelc\_1 X2)\wedge(m2\_finseq\_1 X2 k5\_numbers))\wedge(X0 = k8\_modelc\_1 X1 X2)))) \quad (16)$$

**Theorem 1**

$$\forall X0.((v1\_modelc\_1 X0) \wedge (m2\_finseq\_1 X0 k5\_numbers)) \Rightarrow (v4\_modelc\_1 X0) \Rightarrow (k1\_funct\_1 X0 np\_1 = np\_1)$$