

l26_xreal_0
(TMdCV2GCBK3TRsrpC1Laz3D9SPK7Uoc9cdd)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k11_arytm_3 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow(r1_xxreal_0 X0 X0) \quad (1)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (2)$$

Assume the following.

$$k11_arytm_3 = k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(\forall X2.(v1_xreal_0 X2)\Rightarrow((r1_xxreal_0 X0 X1)\Rightarrow(r1_xxreal_0 (k2_xcmplx_0 X0 X2) (k2_xcmplx_0 X1 X2)))))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow(((r1_xxreal_0 X0 X1)\wedge(r1_xxreal_0 X1 X0))\Rightarrow(X0 = X1))) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(k4_xcmplx_0 (k4_xcmplx_0 X0) = X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k6_xcmplx_0 X0 X1 = k2_xcmplx_0 X0 (k4_xcmplx_0 X1))) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow ((X1 = k4_xcmplx_0 X0) \Leftrightarrow (k2_xcmplx_0 X0 X1 = k6_numbers))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow ((r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (14)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 (k4_xcmplx_0 X1) (k4_xcmplx_0 X0))))$$