

l27_monoid_0

(TMXVSzFooiAof5i5tgd6FdprYZtUhAjP1Rj)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $v5_group_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k6_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(l3_algstr_0 X0) \Rightarrow ((v5_group_1 X0) \Leftrightarrow (\forall X1.(m1_subset_1 \\ X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ X0)) \Rightarrow (k6_algstr_0 X0 X1 X2 = k6_algstr_0 X0 X2 X1)))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow ((v5_group_1 \\ X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k6_algstr_0 X0 X1 X2 = k6_algstr_0 \\ X0 X2 X1)))) \end{aligned}$$