

# l27\_sincos10 (TMFwcWibhxwvoHn- pzArTEc3JFYJV1B2mdmA)

October 27, 2020

Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_rcomp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_real\_1 : \iota \Rightarrow \iota$  be given. Let  $k10\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k32\_sin\_cos : \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k3\_rcomp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k4\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (r1\_tarski (k4\_xxreal\_1 X0 X1) (k2\_xxreal\_1 X0 X1))) \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \quad (3)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (k3\_rcomp\_1 X0 X1 = k2\_xxreal\_1 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (k2\_rcomp\_1 X0 X1 = k4\_xxreal\_1 X0 X1) \quad (6)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0)\wedge(v1\_xxreal\_0 X0) \quad (7)$$

Assume the following.

$$m1\_subset\_1 k32\_sin\_cos k1\_numbers \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(m1\_subset\_1 (k1\_real\_1 X0) k1\_numbers) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers)\wedge(v1\_xreal\_0 X1))\Rightarrow(m1\_subset\_1 (k10\_real\_1 X0 X1) k1\_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(v1\_xxreal\_0 X0) \quad (11)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (12)$$

**Theorem 1**

$$r1\_tarski (k2\_rcomp\_1 (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_2)) k6\_numbers) (k3\_rcomp\_1 (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_2)) k6\_numbers)$$