

l27_sprect_5 (TMYLSMvFw- maSU1mZZy3EoR8AerSgR4erSR3)

October 27, 2020

Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_6 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_topreal1 : \iota \Rightarrow o$ be given. Let $v2_topreal1 : \iota \Rightarrow o$ be given. Let $v1_goboard5 : \iota \Rightarrow o$ be given. Let $v2_goboard5 : \iota \Rightarrow o$ be given. Let $v1_sprect_2 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k20_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k3_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k21_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k24_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0 X0) \wedge ((\neg v3_funct_1 X0) \wedge ((v1_finseq_6 \\ & X0 (u1_struct_0 (k15_euclid np_2))) \wedge ((v1_topreal1 X0) \wedge ((v2_topreal1 \\ & X0) \wedge ((v1_goboard5 X0) \wedge ((v2_goboard5 X0) \wedge ((v1_sprect_2 X0) \wedge \\ & (m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))))))))) \Rightarrow (\\ & \neg(k7_partfun1 (u1_struct_0 (k15_euclid np_2)) X0 np_1 = k20_pscomp_1 \\ & (k3_topreal1 np_2 X0)) \wedge (r1_xxreal_0 (k4_finseq_4 X0 (k24_pscomp_1 \\ & (k3_topreal1 np_2 X0))) (k4_finseq_4 X0 (k21_pscomp_1 (k3_topreal1 \\ & np_2 X0)))))) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v3_funct_1 X0) \wedge ((\neg v1_xboole_0 X0) \wedge ((v1_finseq_6 \\ & X0 (u1_struct_0 (k15_euclid np_2))) \wedge ((v1_topreal1 X0) \wedge ((v2_topreal1 \\ & X0) \wedge ((v1_goboard5 X0) \wedge ((v2_goboard5 X0) \wedge ((v1_sprect_2 X0) \wedge \\ & (m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))))))))) \Rightarrow (\\ & \neg(k7_partfun1 (u1_struct_0 (k15_euclid np_2)) X0 np_1 = k20_pscomp_1 \\ & (k3_topreal1 np_2 X0)) \wedge (r1_xxreal_0 (k4_finseq_4 X0 (k18_pscomp_1 \\ & (k3_topreal1 np_2 X0))) (k4_finseq_4 X0 (k24_pscomp_1 (k3_topreal1 \\ & np_2 X0)))))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (\\ & (v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 \\ & X0))) \Rightarrow (m1_subset_1 (k4_finseq_4 X0 X1) k5_numbers) \end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \tag{9}$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \tag{10}$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (v1_xxreal_0 X0) \tag{11}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v3_funct_1 X0) \wedge ((\neg v1_xboole_0 X0) \wedge ((v1_finseq_6 \\ & X0 (u1_struct_0 (k15_euclid np_2))) \wedge ((v1_topreal1 X0) \wedge ((v2_topreal1 \\ & X0) \wedge ((v1_goboard5 X0) \wedge ((v2_goboard5 X0) \wedge ((v1_sprect_2 X0) \wedge \\ & (m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))))))))) \Rightarrow (\\ & \neg(k7_partfun1 (u1_struct_0 (k15_euclid np_2)) X0 np_1 = k20_pscomp_1 \\ & (k3_topreal1 np_2 X0) \wedge (r1_xxreal_0 (k4_finseq_4 X0 (k18_pscomp_1 \\ & (k3_topreal1 np_2 X0))) (k4_finseq_4 X0 (k21_pscomp_1 (k3_topreal1 \\ & np_2 X0)))))) \end{aligned}$$