

## l28\_qc\_lang4

(TMJVTh4HumhuV4fVEX1bGVJAJJYy3vy3jNj)

October 27, 2020

Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k10\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2))) \Rightarrow ((X2 = k10\_finseq\_1 X0 X1) \Leftrightarrow ((k3\_finseq\_1 X2 = np\_2) \wedge ((k1\_funct\_1 X2 np\_1 = X0) \wedge (k1\_funct\_1 X2 np\_2 = X1)))) \quad (1)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 (k10\_finseq\_1 X0 X1)) \wedge (v1\_funct\_1 (k10\_finseq\_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. v1\_finseq\_1 (k10\_finseq\_1 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (5)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1. (m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow ((X1 = k3\_finseq\_1 X0) \Leftrightarrow (k2\_finseq\_1 X1 = k9\_xtuple\_0 X0))) \quad (6)$$

**Theorem 1**

$$\forall X0.\forall X1.k4\_finseq-1 (k10\_finseq-1 X0 X1) = k2\_finseq-1$$

*np-2*