

## l29\_seqm\_3

(TMdBfHcXjpPtqjK37c4UJGrWpLLJ4tdxD36)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_valued\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1\_funct\_1 X3) \wedge \\ & ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1)))) \Rightarrow (r1\_tarski X1 X2) \Rightarrow (((X1 = k1\_xboole\_0) \wedge (X0 \neq k1\_xboole\_0)) \vee \\ & ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X2) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \tag{2}$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((v1\_funct\_1 X1)\wedge((v1\_funct\_2 X1 k5\_numbers X0)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0))))))\wedge(v7\_ordinal1 X2))\Rightarrow(k8\_nat\_1 X0 X1 X2 = k1\_funct\_1 X1 X2) \quad (5)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0)\wedge((v1\_funct\_2 X0 k5\_numbers k1\_numbers)\wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers))))))\Rightarrow \\ & ((v5\_valued\_0 X0)\Leftrightarrow(\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers)\Rightarrow \\ & (\neg r1\_xreal\_0 (k8\_nat\_1 k1\_numbers X0 (k2\_nat\_1 X1 np\_1)) (k8\_nat\_1 \\ & k1\_numbers X0 X1)))) \quad (8) \end{aligned}$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(v7\_ordinal1 X1))\Rightarrow(v7\_ordinal1 (k2\_xcmplx\_0 X0 X1)) \quad (10)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_nat\_1 X1 X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(v1\_xboole\_0 X1)) \quad (14)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0)\wedge((v1\_funct\_2 X0 k5\_numbers k5\_numbers)\wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k5\_numbers))))))\Rightarrow \\ & ((v5\_valued\_0 X0)\Leftrightarrow(\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers)\Rightarrow \\ & (\neg r1\_xreal\_0 (k8\_nat\_1 k5\_numbers X0 (k2\_nat\_1 X1 np\_1)) (k8\_nat\_1 \\ & k5\_numbers X0 X1)))) \end{aligned}$$