

# l2\_complex1

(TMdmeAUgethAtvHx9nhzec2ycaRCJFXdeF4)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_square\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k3\_xcmplx\_0 X0 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 k6\_numbers X1)) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k2\_xcmplx\_0 X0 X1)))) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xreal\_0 (k3\_square\_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k3\_square\_1 X0 = k3\_xcmplx\_0 X0 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (5)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k2\_xcmplx\_0 (k3\_square\_1 X0) (k3\_square\_1 X1))))$$