

# l2\_convex4 (TMcdCoZoB- HEi1Sv9HmSmaFnjMs7znY9dQdv)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $k9\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k13\_pre\_poly : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (r1\_tarski (k13\_pre\_poly X0) (k9\_xtuple\_0 X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\neg v1\_xboole\_0 k2\_numbers \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X1) \wedge (m1\_funct\_2 X2 X0 X1)) \Rightarrow (\forall X3. (m2\_funct\_2 X3 X0 X1 X2) \Rightarrow ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(\neg v1\_xboole\_0 X1)\Rightarrow(m1\_funct\_2 (k9\_funct\_2 X0 X1) X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow(m1\_subset\_1 (k1\_relset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow(X2 \in X1)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (10)$$

**Theorem 1**

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge(l2\_algstr\_0 X0))\Rightarrow(\forall X1.(m2\_funct\_2 X1 (u1\_struct\_0 X0) k2\_numbers (k9\_funct\_2 (u1\_struct\_0 X0) k2\_numbers))\Rightarrow(r1\_tarski (k13\_pre\_poly X1) (u1\_struct\_0 X0)))$$