

l2_euclid_8 (TMZHZWwrLtredDnLE- hHJM9gBSzHQ9H93QdY)

October 27, 2020

Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k9_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_euclid_8 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k24_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1_xreal_0 X1) \Rightarrow (\forall X2. ((v1_relat_1 \\ & X2) \wedge ((v1_funct_1 X2) \wedge ((v3_valued_0 X2) \wedge (v1_finseq_1 X2)))) \Rightarrow \\ & (k1_seq_1 (k24_valued_1 X2 X1) X0 = k11_binop_2 X1 (k1_seq_1 X2 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. (m2_finseq_2 X0 k1_numbers (k1_euclid np_3)) \Rightarrow (X0 = k1_euclid_8 (k1_seq_1 X0 np_1) (k1_seq_1 X0 np_2) (k1_seq_1 X0 np_3)) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1\ X0)\wedge((m1_subset_1\ X1\ (k1_euclid\ X0))\wedge(v1_xreal_0\ X2)))\Rightarrow(k9_euclid\ X0\ X1\ X2 = k24_valued_1\ X1\ X2) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$v6_membered\ k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2\ X1\ X0)\Rightarrow(\forall X2.(m2_finseq_2\ X2\ X0\ X1)\Rightarrow(m2_finseq_1\ X2\ X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1\ X1\ X0)\Rightarrow((v1_funct_1\ X1)\wedge((v1_finseq_1\ X1)\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ (k2_zfmisc_1\ k5_numbers\ X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1\ X1\ X0)\Rightarrow((v1_relat_1\ X1)\wedge((v1_funct_1\ X1)\wedge(v1_finseq_1\ X1))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1\ X0)\wedge((m1_subset_1\ X1\ (k1_euclid\ X0))\wedge(v1_xreal_0\ X2)))\Rightarrow(m2_finseq_2\ (k9_euclid\ X0\ X1\ X2)\ k1_numbers\ (k1_euclid\ X0)) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(m1_finseq_2\ (k1_euclid\ X0)\ k1_numbers) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1\ X1\ X0)\Rightarrow(v5_relat_1\ X1\ X0) \quad (14)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge(v5_relat_1\ X0\ k1_numbers))\Rightarrow((v1_relat_1\ X0)\wedge(v3_valued_0\ X0)) \quad (15)$$

Assume the following.

$$\forall X0.(v6_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v7_ordinal1\ X1)) \quad (16)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_2 X0 k1_numbers (k1_euclid np_3)) \Rightarrow (\forall X1. \\ & (v1_xreal_0 X1) \Rightarrow (k9_euclid np_3 X0 X1 = k1_euclid_8 (k11_binop_2 \\ & X1 (k1_seq_1 X0 np_1)) (k11_binop_2 X1 (k1_seq_1 X0 np_2)) (k11_binop_2 \\ & X1 (k1_seq_1 X0 np_3)))) \end{aligned}$$