

l2_jordan5a
(TMbhQWzXFL1SxjfkMPppuR5AtJAnhx941fh)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k14_euclid : \iota \Rightarrow \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (u1_struct_0 (k14_euclid X0) = u1_struct_0 (k15_euclid X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (u1_struct_0 (k15_euclid X0) = k1_euclid X0) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (4)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow ((u1_struct_0 (k14_euclid X0) = k1_euclid X0) \wedge (u1_struct_0 (k15_euclid X0) = k1_euclid X0))$$