

## l2\_lexbfs

(TMG992TWjUe8fr1vWwWUaYy4dKWW7DJmvoe)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v2\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_funct\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relat\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1\_relat\_1 X2) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (r1\_tarski (k5\_relat\_1 X2 X0) (k5\_relat\_1 X2 X1))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((v2\_funct\_1 X1) \Rightarrow (v2\_funct\_1 (k5\_relat\_1 X1 X0))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Leftrightarrow ((r1\_tarski (k9\_xtuple\_0 X0) (k9\_xtuple\_0 X1)) \wedge (\forall X2. (X2 \in k9\_xtuple\_0 X0) \Rightarrow (k1\_funct\_1 X0 X2 = k1\_funct\_1 X1 X2)))))) \quad (3)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Rightarrow (\forall X1. (v1\_relat\_1 X1) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (r1\_tarski (k2\_relat\_1 X0) (k2\_relat\_1 X1)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow ((v1\_relat\_1 (k5\_relat\_1 X0 X1)) \wedge (v1\_funct\_1 (k5\_relat\_1 X0 X1))) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X0) \Rightarrow (v1\_relat\_1 (k5\_relat\_1 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow ((v1\_relat\_1 (k2\_funct\_1 X0)) \wedge (v1\_funct\_1 (k2\_funct\_1 X0))) \quad (7)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow ((v2\_funct\_1 X0) \Rightarrow (k2\_funct\_1 X0 = k2\_relat\_1 X0)) \quad (8)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v2\_funct\_1 X0))) \Rightarrow \\ & (\forall X1. \forall X2. (r1\_tarSKI X1 X2) \Rightarrow (\forall X3. (X3 \in k9\_xtuple\_0 \\ & (k2\_funct\_1 (k5\_relat\_1 X0 X1))) \Rightarrow (k1\_funct\_1 (k2\_funct\_1 (k5\_relat\_1 \\ & X0 X1)) X3 = k1\_funct\_1 (k2\_funct\_1 (k5\_relat\_1 X0 X2)) X3))) \end{aligned}$$