

l2_mboolean (TM- Liv7LXroLTLqhXMHCqRG928ogfBm7UQRi)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X1 \in k9_xtuple_0 X0) \Rightarrow (k9_xtuple_0 X0 = k9_xtuple_0 (k1_funct_4 \\ & X0 (k16_funcop_1 X1 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & (v1_partfun1 X1 X0) \Leftrightarrow (k1_relset_1 X0 X1 = X0)) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_relat_1 X3) \wedge \\ & ((v4_relat_1 X3 X1) \wedge ((v1_funct_1 X3) \wedge (v1_partfun1 X3 X1)))) \Rightarrow \\ & ((X0 \in X1) \Rightarrow (k9_xtuple_0 (k1_funct_4 X3 (k16_funcop_1 X0 X2)) = X1)) \end{aligned}$$