

l2_pepin

(TMT7jiT7qAEeFUtuudeaTxj8XEbvJNqAmi6)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg r1_xxreal_0 (k2_newton np_2 X0) X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xxreal_0 k6_numbers X0) \quad (3)$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge (v7_ordinal1 X0)) \Rightarrow (\neg (X0 \neq np_1) \wedge (r1_xxreal_0 X0 np_1)) \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((\neg r1_xxreal_0 np_1 X0) \Rightarrow (X0 = k6_numbers)) \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_newton X0 X1 = k1_newton X0 X1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k1_newton X0 X1)) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (12)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\neg r1_xxreal_0 (k2_newton np_2 (k2_newton np_2 X0)) np_1)$$