

l2_radix_3

(TMZs1t8LYkU3HJQ5GPYrdQeLXFrnrhyRG3)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_radix_1 : \iota \Rightarrow \iota$ be given. Let $k7_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k3_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\\ & (r1_xxreal_0 np_1 (k7_nat_d X0 X1)) \vee (r1_xxreal_0 np_1 (k6_xcmplx_0 \\ & X0 X1))) \Rightarrow (k7_nat_d X0 X1 = k6_xcmplx_0 X0 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k7_nat_d (k2_xcmplx_0 X0 X1) X1 = X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 np_1 X0) \Rightarrow (r1_xxreal_0 np_1 (k1_newton X0 X1)))) \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$r1_xxreal_0 np_1 np_2 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((r1_xreal_0 np_1 X0) \Rightarrow (k1_radix_1 X0 = k2_xcmplx_0 (k1_radix_1 (k7_nat_d X0 np_1)) (k1_radix_1 (k7_nat_d X0 np_1)))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (k3_power X0 X1 = k1_newton X0 X1) \quad (9)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (m1_subset_1 (k7_nat_d X0 X1) k5_numbers) \quad (11)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (m1_subset_1 (k1_radix_1 X0) k5_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k1_radix_1 X0 = k3_power np_2 X0) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (14)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (15)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (16)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((r1_xreal_0 np_1 X0) \Rightarrow (k6_xcmplx_0 (k1_radix_1 X0) (k1_radix_1 (k7_nat_d X0 np_1)) = k1_radix_1 (k7_nat_d X0 np_1)))$$