

l2_rewrite1

(TMXyz9KoaCqic7xRZnqdVBkj1pnLVeYrFC8)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ & ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 k6_numbers = X0) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\\ & \neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$(m2_subset_1\ np_0\ k1_numbers\ k5_numbers) \wedge ((m1_subset_1\ np_0\ k5_numbers) \wedge (m1_subset_1\ np_0\ k1_numbers)) \quad (6)$$

Assume the following.

$$v1_xboole_0\ np_0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0) \wedge ((v1_funct_1\ X0) \wedge (v1_finseq_1\ X0))) \Rightarrow (k4_finseq_1\ X0 = k9_xtuple_0\ X0) \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0) \wedge ((v1_funct_1\ X0) \wedge (v1_finseq_1\ X0))) \Rightarrow (k3_finseq_1\ X0 = k1_card_1\ X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1\ X0\ k5_numbers) \wedge (v7_ordinal1\ X1)) \Rightarrow (k2_nat_1\ X0\ X1 = k2_xcmplx_0\ X0\ X1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1\ X1) \wedge (v4_relat_1\ X1\ X0)) \Rightarrow (k1_relset_1\ X0\ X1 = k9_xtuple_0\ X1) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0) \wedge (m1_subset_1\ X1\ k5_numbers)) \Rightarrow (k1_nat_1\ X0\ X1 = k2_xcmplx_0\ X0\ X1) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1\ X0\ k5_numbers) \wedge (v7_ordinal1\ X1)) \Rightarrow (k2_nat_1\ X0\ X1 = k2_nat_1\ X1\ X0) \quad (15)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1) \Rightarrow (v7_ordinal1\ X0) \quad (16)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0) \Rightarrow (v7_ordinal1\ X0) \quad (17)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge \\ (v1_finseq_1 X0)))) \end{aligned} \quad (18)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (20)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1.(v7_ordinal1 X1) \Rightarrow ((X1 \in k4_finseq_1 X0) \Rightarrow ((r1_xxreal_0 \\ X1 (k3_finseq_1 X0)) \wedge ((r1_xxreal_0 (k2_nat_1 k6_numbers np_1) \\ X1) \wedge (\neg r1_xxreal_0 X1 k6_numbers)))))) \end{aligned}$$