

# l2\_subset\_1 (TMXmjPVNtKXLzUQRFPTX- iqya2yb2zSmyVvi)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \neg v1\_xboole\_0 (k1\_zfmisc\_1 X0) \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1\_zfmisc\_1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (r1\_tarski X2 X0)) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \Rightarrow ((m1\_subset\_1 X1 X0) \Leftrightarrow (X1 \in X0))) \wedge ((v1\_xboole\_0 X0) \Rightarrow ((m1\_subset\_1 X1 X0) \Leftrightarrow (v1\_xboole\_0 X1))) \tag{4}$$

**Theorem 1**

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (\forall X2. (X2 \in X1) \Rightarrow (X2 \in X0))$$