

l2_valued_1 (TM- SaSe1GmMpbNBbVKhUk8mojMYHQwMcmCCF)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k2_finseq_1 X0 = k9_subset_1 k5_numbers (k2_finseq_1 X1) (k2_finseq_1 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m1_subset_1 (k4_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow ((v1_finseq_1 X0) \Leftrightarrow (\exists X1.(v7_ordinal1 X1) \wedge (k9_xtuple_0 X0 = k2_finseq_1 X1))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\begin{aligned} &\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow \\ &\quad (\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 \\ &X1)))\Rightarrow(\forall X2.((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((k9_xtuple_0 \\ &X2 = k9_subset_1 k5_numbers (k4_finseq_1 X0) (k4_finseq_1 X1))\Rightarrow \\ &\quad ((v1_relat_1 X2)\wedge((v1_funct_1 X2)\wedge(v1_finseq_1 X2)))))) \end{aligned}$$