

l2_waybel14

(TMX2ary7xCi9D4JMpCzmUyEmxt7AVptMmG8)

October 27, 2020

Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow ((r1_xboole_0 X1 X2) \Leftrightarrow (r1_tarski \\ X1 (k3_subset_1 X0 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Rightarrow (r1_xboole_0 X1 X0) \tag{2}$$

Theorem 1

$$\begin{aligned} \forall X0. (l1_struct_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow ((r1_tarski X1 (k3_subset_1 (u1_struct_0 \\ X0) X2)) \Leftrightarrow (r1_tarski X2 (k3_subset_1 (u1_struct_0 X0) X1)))))) \end{aligned}$$