

l33_o_ring_1

(TMEhDE2qj2wWMgFKaUy9tNvLyoAwt9rcjeH)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l6_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_o_ring_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v11_o_ring_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v10_o_ring_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l5_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l6_algstr_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow ((v1_o_ring_1 X1 X0) \Rightarrow (v10_o_ring_1 \\ & (k12_finseq_1 (u1_struct_0 X0) X1) X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l6_algstr_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m2_finseq_1 X2 \\ & (u1_struct_0 X0)) \Rightarrow ((X2 = k12_finseq_1 (u1_struct_0 X0) X1) \Leftrightarrow ((\\ & k3_finseq_1 X2 = np_1) \wedge (k7_partfun1 (u1_struct_0 X0) X2 np_1 = \\ & X1)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \tag{3}$$

Assume the following.

$$\forall X0.(l6_algstr_0 X0) \Rightarrow ((l2_algstr_0 X0) \wedge (l5_algstr_0 X0)) \tag{4}$$

Assume the following.

$$\forall X0.(l2_struct_0 X0) \Rightarrow (l1_struct_0 X0) \tag{5}$$

Assume the following.

$$\forall X0.(l2_algstr_0 X0) \Rightarrow ((l2_struct_0 X0) \wedge (l1_algstr_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (7)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l6_algstr_0 X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow ((v11_oring_1 X1 X0) \Leftrightarrow (\exists X2. (m2_finseq_1 X2 (u1_struct_0 X0)) \wedge ((v10_oring_1 X2 X0) \wedge (X1 = k7_partfun1 (u1_struct_0 X0) X2 (k3_finseq_1 X2))))))) \quad (8)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l6_algstr_0 X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow ((v1_oring_1 X1 X0) \Rightarrow (v11_oring_1 X1 X0)))$$