

l33_toprealb

(TMJ3MLZiJyVpfsd1qzRvFvJxvyVj8QoLjt4)

October 27, 2020

Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k8_toprealb : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k3_topreal9 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $k7_toprealb : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(v1_xreal_0 \\ & X1) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid X0))) \Rightarrow \\ & (u1_struct_0 (k7_toprealb X0 X2 X1) = k3_topreal9 X0 X2 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$k4_struct_0 (k15_euclid np_2) = k19_euclid k6_numbers k6_numbers \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (8)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k19_euclid X0 X1) (u1_struct_0 (k15_euclid np_2))) \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k8_toprealb X0 = k7_toprealb X0 (k4_struct_0 (k15_euclid X0)) np_1) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (13)$$

Theorem 1

$$u1_struct_0 (k8_toprealb np_2) = k3_topreal9 np_2 (k19_euclid k6_numbers k6_numbers) np_1$$