

l36_modelc_1

(TMca9ahYEGhwseVQTjaCNiAxEgEP84UpLJ2)

October 27, 2020

Let $v1_modelc_1 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v2_modelc_1 : \iota \Rightarrow o$ be given. Let $v3_modelc_1 : \iota \Rightarrow o$ be given. Let $v4_modelc_1 : \iota \Rightarrow o$ be given. Let $v5_modelc_1 : \iota \Rightarrow o$ be given. Let $v6_modelc_1 : \iota \Rightarrow o$ be given. Let $v7_modelc_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_2 : \iota$ be given. Let $np_3 : \iota$ be given. Let $np_4 : \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow & \\ (v2_modelc_1 X0) \Rightarrow ((k1_funct_1 X0 np_1 \neq k6_numbers) \wedge ((k1_funct_1 & \\ X0 np_1 \neq np_1) \wedge ((k1_funct_1 X0 np_1 \neq np_2) \wedge ((k1_funct_1 X0 & \\ np_1 \neq np_3) \wedge (k1_funct_1 X0 np_1 \neq np_4)))))) & \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow & \\ (v7_modelc_1 X0) \Rightarrow (k1_funct_1 X0 np_1 = np_4) & \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow & \\ (v6_modelc_1 X0) \Rightarrow (k1_funct_1 X0 np_1 = np_3) & \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow & \\ (v5_modelc_1 X0) \Rightarrow (k1_funct_1 X0 np_1 = np_2) & \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow & \\ (v4_modelc_1 X0) \Rightarrow (k1_funct_1 X0 np_1 = np_1) & \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow & \\ (v3_modelc_1 X0) \Rightarrow (k1_funct_1 X0 np_1 = k6_numbers) & \end{aligned} \quad (6)$$

Theorem 1

$$\forall X0.((v1_modelc_1 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (v2_modelc_1 X0) \Rightarrow ((\neg v3_modelc_1 X0) \wedge ((\neg v4_modelc_1 X0) \wedge ((\neg v5_modelc_1 X0) \wedge ((\neg v6_modelc_1 X0) \wedge (\neg v7_modelc_1 X0)))))$$