

l36_modelc_2

(TMLLCTKtdX8JdhUTbZasiRfC6TZreHrEgNc)

October 27, 2020

Let $v1_modelc_2 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v3_modelc_2 : \iota \Rightarrow o$ be given. Let $v2_modelc_2 : \iota \Rightarrow o$ be given. Let $v4_modelc_2 : \iota \Rightarrow o$ be given. Let $v5_modelc_2 : \iota \Rightarrow o$ be given. Let $v6_modelc_2 : \iota \Rightarrow o$ be given. Let $v7_modelc_2 : \iota \Rightarrow o$ be given. Let $v8_modelc_2 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $np_5 : \iota$ be given. Let $np_4 : \iota$ be given. Let $np_3 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_zfmisc_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\neg v1_xboole_0 \ np_5 \tag{1}$$

Assume the following.

$$\neg v1_xboole_0 \ np_4 \tag{2}$$

Assume the following.

$$\neg v1_xboole_0 \ np_3 \tag{3}$$

Assume the following.

$$\neg v1_xboole_0 \ np_2 \tag{4}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_2 \ X0) \wedge (m2_finseq_1 \ X0 \ k5_numbers)) \Rightarrow (\\ (v2_modelc_2 \ X0) \Rightarrow ((k1_funct_1 \ X0 \ np_1 \neq k6_numbers) \wedge ((k1_funct_1 \\ X0 \ np_1 \neq np_1) \wedge ((k1_funct_1 \ X0 \ np_1 \neq np_2) \wedge ((k1_funct_1 \ X0 \\ np_1 \neq np_3) \wedge ((k1_funct_1 \ X0 \ np_1 \neq np_4) \wedge (k1_funct_1 \ X0 \ np_1 \neq \\ np_5))))))))) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_modelc_2 \ X0) \wedge (m2_finseq_1 \ X0 \ k5_numbers)) \Rightarrow (\\ (v8_modelc_2 \ X0) \Rightarrow (k1_funct_1 \ X0 \ np_1 = np_5)) \end{aligned} \tag{7}$$

Assume the following.

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v7_modelc_2 X0)\Rightarrow(k1_funct_1 X0 np_1 = np_4)) \quad (8)$$

Assume the following.

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v6_modelc_2 X0)\Rightarrow(k1_funct_1 X0 np_1 = np_3)) \quad (9)$$

Assume the following.

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v5_modelc_2 X0)\Rightarrow(k1_funct_1 X0 np_1 = np_2)) \quad (10)$$

Assume the following.

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v4_modelc_2 X0)\Rightarrow(k1_funct_1 X0 np_1 = np_1)) \quad (11)$$

Assume the following.

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v3_modelc_2 X0)\Rightarrow(k1_funct_1 X0 np_1 = k6_numbers)) \quad (12)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (13)$$

Assume the following.

$$\forall X0.(v3_card_1 X0 np_1)\Rightarrow((\neg v1_xboole_0 X0)\wedge(v1_zfmisc_1 X0)) \quad (14)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v3_card_1 X0 k1_xboole_0) \quad (15)$$

Theorem 1

$$\forall X0.((v1_modelc_2 X0)\wedge(m2_finseq_1 X0 k5_numbers))\Rightarrow((v3_modelc_2 X0)\Rightarrow((\neg v2_modelc_2 X0)\wedge((\neg v4_modelc_2 X0)\wedge((\neg v5_modelc_2 X0)\wedge((\neg v6_modelc_2 X0)\wedge((\neg v7_modelc_2 X0)\wedge(\neg v8_modelc_2 X0))))))))$$