

l37_matrix13

(TMEtC8fmyaD8ZqMsgrrvVqugKkkSAqRqQZk)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow (((k10_xtuple_0 X0 = k10_xtuple_0 X1) \wedge ((k3_finseq_1 X0 = \\ & k3_finseq_1 X1) \wedge (v2_funct_1 X1))) \Rightarrow (v2_funct_1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (5)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Rightarrow(m2_finseq_1 X2 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(v7_ordinal1 X0)\Rightarrow(m1_finseq_2 (k4_finseq_2 X0 X1) X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1 X1 X0)\Leftrightarrow(k1_card_1 X1 = X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(v7_ordinal1 X1))\Rightarrow(\forall X2.(m1_subset_1 X2 (k4_finseq_2 X1 X0))\Rightarrow(v3_card_1 X2 X1)) \quad (11)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(m2_finseq_2 X1 k5_numbers (k4_finseq_2 X0 k5_numbers))\Rightarrow(\forall X2.(m2_finseq_2 X2 k5_numbers (k4_finseq_2 X0 k5_numbers))\Rightarrow(((k10_xtuple_0 X1 = k10_xtuple_0 X2)\wedge(v2_funct_1 X1))\Rightarrow(v2_funct_1 X2))))$$