

l37_random_1

(TMPghSiPV9NN4MkgHpXe5v4zLTiTqBdx6p)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $k56_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $k54_valued_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow ((\forall X2. (X2 \in k1_relset_1 \\ X0 X1) \Rightarrow (r1_xxreal_0 k6_numbers (k1_seq_1 X1 X2))) \Rightarrow (v6_supinf_2 \\ X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k17_complex1 X0)) \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((v1_membered X1) \wedge ((v1_funct_1 \\ X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow (k56_valued_1 \\ X0 X1 X2 = k54_valued_1 X2)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow(v1_xcmplx_0 (k1_funct_1 X0 X1)) \quad (6)$$

Assume the following.

$$v3_membered k1_numbers \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_membered X1)\wedge((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))\Rightarrow((v1_funct_1 (k56_valued_1 X0 X1 X2))\wedge(m1_subset_1 (k56_valued_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v3_valued_0 X1))))\Rightarrow((X1 = k54_valued_1 X0)\Leftrightarrow((k9_xtuple_0 X1 = k9_xtuple_0 X0)\wedge(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k1_funct_1 X1 X2 = k17_complex1 (k1_funct_1 X0 X2)))))) \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v3_valued_0 X0))\Rightarrow((v1_relat_1 X0)\wedge(v1_valued_0 X0)) \quad (10)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(v1_membered X0) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (14)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow(v6_supinf_2 (k56_valued_1 X0 k1_numbers X1)))$$