

l39_cqc_the3

(TMYkXSxC8dEwSt5YX7pvBDVp92P5u9C9knu)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r3_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r4_cqc_the3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_cqc_the3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (r4_cqc_the3 X0 (k7_domain_1 (k3_cqc_lang \\ X0) X1 X2) (k6_domain_1 (k3_cqc_lang X0) (k7_cqc_lang X0 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (((r4_cqc_the3 X0 X1 X2) \wedge (r2_cqc_the3 X0 X3 \\ X1)) \Rightarrow (r2_cqc_the3 X0 X3 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow ((r2_cqc_the3 X0 X2 (k6_domain_1 (k3_cqc_lang \\ X0) X1)) \Leftrightarrow (r3_cqc_the1 X0 X2 X1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge((m1_subset_1 X1 X0)\wedge(m1_subset_1 X2 X0)))\Rightarrow(k7_domain_1 X0 X1 X2 = k2_tarski X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(k6_domain_1 X0 X1 = k1_tarski X1) \quad (6)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\neg v1_xboole_0 (k3_cqc_lang X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge((m1_subset_1 X1 X0)\wedge(m1_subset_1 X2 X0)))\Rightarrow(m1_subset_1 (k7_domain_1 X0 X1 X2) (k1_zfmisc_1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0)\wedge((m1_subset_1 X1 (k3_cqc_lang X0))\wedge(m1_subset_1 X2 (k3_cqc_lang X0))))\Rightarrow(m2_subset_1 (k7_cqc_lang X0 X1 X2) (k9_qc_lang1 X0) (k3_cqc_lang X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(m1_subset_1 (k6_domain_1 X0 X1) (k1_zfmisc_1 X0)) \quad (10)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(m1_subset_1 (k3_cqc_lang X0) (k1_zfmisc_1 (k9_qc_lang1 X0))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k2_tarski X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 = X0)\vee(X3 = X1))) \quad (12)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k3_cqc_lang X0)))\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k3_cqc_lang X0)))\Rightarrow((r2_cqc_the3 X0 X1 X2)\Leftrightarrow(\forall X3.(m2_subset_1 X3 (k9_qc_lang1 X0) (k3_cqc_lang X0))\Rightarrow((X3 \in X2)\Rightarrow(r3_cqc_the1 X0 X1 X3)))))) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (14)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X3.(m1_subset_1\ X3\ (k1_zfmisc_1 \\ (k3_cqc_lang\ X0))) \Rightarrow (((r3_cqc_the1\ X0\ X3\ X1) \wedge (r3_cqc_the1\ X0\ X3 \\ X2)) \Rightarrow (r3_cqc_the1\ X0\ X3\ (k7_cqc_lang\ X0\ X1\ X2)))))) \end{aligned}$$