

## l3\_lattice8

(TMb4zXJ6R3HV2NVGB4R4KUHP6YpAwiR3HmM)

October 27, 2020

Let  $v1\_abian : \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_int\_1 X0) \Rightarrow ((v1\_abian X0) \Leftrightarrow (\neg v1\_abian (k6\_xcmplx\_0 X0 np\_1))) \quad (1)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (2)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_int\_1 X0) \wedge (v1\_abian X0)) \wedge ((v1\_int\_1 X1) \wedge (v1\_abian X1))) \Rightarrow (v1\_abian (k6\_xcmplx\_0 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_int\_1 X0) \quad (6)$$

**Theorem 1**  $\neg v1\_abian np\_1$ .