

l3_modelc_3

(TMQKjH6MmQF4tptu8YbLvf4fWeGjPmLeqocg)

October 27, 2020

Let $v1_modelc_2 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((k9_xtuple_0 \\ & X1 = k1_tarski X0) \Rightarrow (k10_xtuple_0 X1 = k1_tarski (k1_funct_1 X1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$(k2_finseq_1 np_1 = k1_tarski np_1) \wedge (k2_finseq_1 np_2 = k2_tarski np_1 np_2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (k2_relset_1 X0 X1 = k10_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (6)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k5_finseq_1 X0)) \wedge (v1_funct_1 (k5_finseq_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k9_xtuple_0 X1 = k2_finseq_1 np_1) \wedge (k1_funct_1 X1 np_1 = X0))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0) \Rightarrow (v5_relat_1 X1 X0) \quad (12)$$

Theorem 1

$$\forall X0.((v1_modelc_2 X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow ((k1_funct_1 (k12_finseq_1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k5_numbers)) X0) np_1 = X0) \wedge (k2_relset_1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k5_numbers)) (k12_finseq_1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k5_numbers)) X0) = k1_tarski X0))$$